

HAMADRYAD

Accepted Manuscript



Title A regurgitation event in Indian Ratsnake *Ptyas mucosa* (LINNAEUS, 1758) provides insights into its diet

Authors Kirubakaran Samson¹, Anbazhagan Abinesh², C.S. Vishnu^{3*}, R. Sagadevan⁴

¹Department of Rural Development Science, Arul Anandar College, Karumathur, Madurai-625514.

²409/155 Lakshmi Nanjan Nivas, Stanley Park, Coonoor- The Nilgiris, Tamil Nadu, 643105.

³Wildlife Institute of India, Chandrabani, Dehradun-248001.

⁴2/236 Thottam, Vedarpuliyangulam, Madurai – 625006.

*Corresponding author: vishnusreedharannair@gmail.com

Date submitted 07/03/2023

Date accepted 30/07/2023

Available online

Citation Samson, K., Abinesh, A., Vishnu, C.S. and Sagadevan, R. (2023) A regurgitation event in Indian Ratsnake *Ptyas mucosa* (LINNAEUS, 1758) provides insights into its diet. *Hamadryad*. In Press.

IMPORTANT This current version of the accepted manuscript is a submission that has been peer reviewed and accepted for publication by *Hamadryad*. This version will have minor differences from the final publication. During the article production process, the text and other elements of this article are likely to change.

When the final article is available, the ‘Accepted Manuscript’ version will be removed and replaced by the final article. The date the article was first made available online will be carried over, but the year of publication will correspond to the volume/issue of inclusion.

An accepted manuscript at *Hamadryad* can be cited in text as “In Press”.

Keywords: Birds, Diet, Eggs, Feeding, Rodents

The prey-predator relationship is a key component that explains the trophic ecology of an environment (Dorresteijn et al. 2015). Functionally, the mechanised prey-predator dynamics conceptualise the trophic cascade, which determines changes in prey abundance and behaviour (Silliman and Angelini 2012). The predatory dichotomy that includes active and ambush predators differ in foraging mode, which intertwines with morphological and behavioural traits through the adaptive syndrome hypothesis (Eckhardt 1979; McLaughlin 1989). The adaptive syndrome can enhance behavioural syndrome, which explains how behaviour changes in different foraging scenarios (Sih et al. 2004). The community profile is the size profile of prey species, and the food profile is the ingested species from the community profile, which is mediated through the prey-predatory dynamics (Griffiths 1975). The optimal foraging theory involves animals balancing energy expenditure with each predation event's success to avoid starvation. This relates to trade-offs in foraging ecology (Higginson and Ruxton 2015). The foraging pattern of predators can also switch depending on prey availability, abundance, and how widely the area is occupied by the prey species (Hirvonen 1999; Higginson and Ruxton 2015). Predators may choose their prey based on factors such as size and predation cost. Among active foragers, there is likely to be higher energy expenditure compared to ambush foragers (Downes 2002; Higginson and Ruxton 2015). This note reports an observation of the stomach contents in the active foraging Indian rat snake (*Ptyas mucosa*). The Indian rat snake is a common, widespread, non-venomous snake found across India in different habitats such as xeric regions, mesic habitats, coastal lines, open fields, deciduous forests, evergreen forests, scrub jungles, and agricultural lands (Whitaker and Captain 2004; Parmar and Tank 2019; Biodiversity portal 2023). The diet of *P. mucosa* varies with age and habitat. Juveniles primarily feed on insects, reptiles, and frogs but shift to consuming mammals, birds, fishes, amphibians, snakes, and other reptiles as they grow (Parmar and Patel 2022).

On 4th October 2018 at 17.00 h, we got a rescue call about a snake's presence inside a house near Thiagarajar School of Management on Avaniyapuram road, Thirupparankundram, Madurai, Tamil Nadu (9.8796°N; 78.0863°E). We rushed to the spot and found an Indian Rat snake (*Ptyas mucosa*) approximately 7 feet in size hanging on the window. We gently rescued it from the house and brought it outside (Figure 1). Instantly, the snake started regurgitating a variety of prey items, including a marbled balloon frog (*Uperodon systoma*). This individual had an olive-yellowish dorsum with a marbled design and a smaller head with a blunt snout. We also spotted an unidentified rodent and a juvenile three-striped palm squirrel (*Funambulus palmarum*) with three stripes on its dorsum. Additionally, we found two bird eggs — one was broken with a partially grown juvenile bird inside. Later identification revealed that the eggs belonged to a white-breasted water hen (*Amaurornis phoenicurus*; Anthal and Sahi 2017) (Figure 1). Subsequently, the snake was safely bagged and released into a nearby plot of agriculture land (9.8840°N; 78.0891°E).

Several prey items of *P. mucosa* have been reported in the literature, including *Eutropis dissimilis*, *Hoplobatrachus tigerinus*, *Euphlyctis cyanophlyctis*, *Fejervarya limnocharis*, and unidentified toads (Wall 1926; Dunn 1935; Minton 1966; Sharma and Vazirani 1977; Whitaker and Captain 2004). Studies have also documented unusual feeding and scavenging events in *P. mucosa*, such as consumption of plastics, onions, male contraceptives, cloth pieces, discarded

socks, and polythene rolls (Sharma et al. 2016; Saha and Chaudhuri 2017; Chaudhuri et al. 2018; Parmar and Patel 2022).

In general, stomach content analyses in snakes have reported details on the diet and prey species that are known to occur in an ecosystem. However, this observation on the regurgitation by this *P. mucosa* individual indicates that the diet can consist of multiple species and diversified prey preferences in *P. mucosa*. This predation strategy may help maintain their metabolic energy requirements, as they are active foragers. Foraging is a basic and primary trade-off activity that limits prey dynamics such as availability, abundance, and spatiotemporal distribution (Higginson and Ruxton 2015). Higginson and Ruxton (2015) explained a relationship between prey statuses to predatory movement in environmental space, stressing the fact that active predators have a higher prey cost. The pressure of food consumption to either overcome or avoid starvation may cause the predator to engage in opportunistic foraging, and preying on what is available, which is seemingly costly, yet a viable option for generalists (Westoby 1978; Rex et al. 2010; Pereira et al. 2016).

The stomach content also gives an idea of the diet of a species, which would be important for the conservation of prey and predators. Regurgitation and aggressive behaviour during handling could be a defence mechanism to allow for a quick escape.

Perez-Ramos et al. (2018) reported regurgitation of three painted wood rats (*Neotoma picta*) by Barbour's montane pit viper (*Mixcoatalus barbouri*) from Mexico. This note reports the diet of *P. mucosa*, which includes different prey species of smaller size, indicating the individual's need to be an active forager in an environment where larger prey is lacking. However, we are aware that *P. mucosa* has a generalist diet and it may or may not consume a wide range of species at a single time. Since we could not find any other related observations for this snake, we conclude this to be significant. More studies on foraging aspects along with stomach content analyses across a spatiotemporal gradient would enhance the understanding of foraging behaviour and ecology of this common colubrid.

Acknowledgement

We would like to acknowledge the Tamil Nadu Forest Department for granting permission to rescue the snake. Additionally, our gratitude goes to Mr. Vivek Sharma for encouraging us to write this observation.

Reference

- Anthal, A. & Sahi, D.N. (2017) A study on clutch and egg characteristics of White-breasted Waterhen, *Amaurornis phoenicurus phoenicurus* in Jammu (J&K), India. *International Journal of Current Research* 9 (1), 4539745400.
- Chaudhuri, A., Sardar, A., Chowdhury, S. & Dutta, A. (2018) *Ptyas mucosa* (Indian Ratsnake). Diet and behavior. *Herpetological Review* 49, 352–353.
- Dorresteijn, I., Schultner, J., Nimmo, D.G., Fischer, J., Hanspach, J., Kuemmerle, T., Kehoe, L. & Ritchie, E.G. (2015) Incorporating anthropogenic effects into trophic ecology: predator–prey interactions in a human-dominated landscape. *Proceedings of the Royal Society B: Biological Sciences* 282(1814), 20151602.
- Downes, S.J. (2002) Size-dependent predation by snakes: selective foraging or differential prey vulnerability?. *Behavioral Ecology* 13(4), 551–560.

- Dunn, E.R. (1935) The Reptiles of China: Natural History of Central Asia, Vol. X, The Reptiles of China. By Clifford H. Pope, *American Museum of Natural History, Science* 82(2126), 303–304.
- Eckhardt, R.C. (1979) The adaptive syndromes of two guilds of insectivorous birds in the Colorado Rocky Mountains. *Ecological Monographs* 49(2), 129–149.
- Griffiths, D. (1975) Prey availability and the food of predators. *Ecology* 56(5), 1209–1214.
- Higginson, A.D. & Ruxton, G. D. (2015) Foraging mode switching: the importance of prey distribution and foraging currency. *Animal Behaviour* 105, 121–137.
- Hirvonen, H. (1999) Shifts in foraging tactics of larval damselflies: effects of prey density. *Oikos* 443–452.
- India Biodiversity Portal. (2023) *Ptyas mucosa*. Available from: <https://indiabiodiversity.org/species/show/257422/> (March 07, 2023)
- Mclaughlin, R.L. (1989) Search modes of birds and lizards: Evidence for alternative movement patterns. *The American Naturalist* 133(5), 654–670.
- Minton, S.A. (1966) A contribution to the herpetology of West Pakistan. *Bulletin of the American Museum of Natural History* 134(2), 27–184.
- Parmar, D.S. & Patel, V.D. (2022) Plastic ingestion by the Indian snakes *Ptyas mucosa* and *Coelognathus helena helena* (Serpentes: Colubridae). *Phyllomedusa: Journal of Herpetology* 21(1), 91–94.
- Parmar, D. & Tank, S. (2019) Herpetofauna of Veer Narmad South Gujarat University, Surat, India. *IRCF Reptiles & Amphibians* 26 (1): 21–34.
- Pereira, S.C., Rodrigues, J.F.M. & Borges-Nojosa, D.M. (2016) Single large or several small?: The influence of prey size on feeding performance of *Philodryas nattereri* (Squamata: Serpentes). *Acta Scientiarum - Biological Sciences* 38, (1), 25–28.
- Perez-Ramos, E., León-Paniagua, L. & Mociño-Deloya, E. (2017) *Mixcocoatlus barbouri* (Barbour's Montane Pitviper) Diet. *Herpetological Review* 48 (3), 678.
- Rex, K., Czaczkes, B.I., Michener, R., Kunz, T.H. & Voigt, C.C. (2010) Specialization and omnivory in diverse mammalian assemblages. *Ecoscience* 17(1), 37–46.
- Saha, A. & Chaudhuri, A. (2017) *Ptyas mucosa* (Indian Rat Snake). Diet / Cannibalism. *Herpetological Review* 48(3), 681.
- Sharma, R.C. & Vazirani, T.G. (1977) Food and feeding habits of some reptiles of Rajasthan. *Records of the Zoological Survey of India* 73(1–4), 77–93.
- Sharma, V., Sayyed, A. & Bhandari, R. (2016) Herbivory and inanimate objects in the diet of the Oriental Ratsnake, *Ptyas mucosa* (Linnaeus 1758). *Reptiles & Amphibians* 23(2), 102–103.
- Sih, A., Bell, A. & Johnson, J.C. (2004) Behavioral syndromes: an ecological and evolutionary overview. *Trends in ecology & evolution* 19(7), 372–378.

- Silliman, B.R. & Angelini, C. (2012) Trophic cascades across diverse plant ecosystems. *Nature education knowledge* 3(10).
- Wall, F. (1926) Snakes collected in Burma in 1925. *Journal of the Bombay Natural History Society* 31(3), 558–566.
- Westoby, M. (1978) What are the biological bases of varied diets?. *The American Naturalist* 112(985), 627–631.
- Whitaker, R. & Captain, A. (2004) *Snakes of India*. Macmillan India Limited, New Delhi, 124.

Accepted
Manuscript



Figure 1. An array of regurgitated meals by *P. mucosa*, showcasing species names from left to right: *Amaurornis phoenicurus*, *Funambulus palmarum*, Unidentified Rodent, *Uperodon systoma*, and White-breasted Waterhen Eggs.