

# HAMADRYAD

## Accepted Manuscript



**Title** First record of intraspecific kleptoparasitism as a foraging strategy in anurans: Observed in the Matheran leaping-frog (*Indirana leithii*) from Maharashtra, India

**Author** Prathamesh Amberkar <sup>1\*</sup>

<sup>1</sup> Ashoka Trust for Research in Ecology and the Environment, Royal Enclave, Srirampura, Jakkur Post, Bangalore - 560064, India.

\*Corresponding author: prathameshsamberkar@gmail.com

**Date submitted** 08/06/2023

**Date accepted** 12/08/2023

**Available online**

**Citation** Amberkar, P. (2023) First record of intraspecific kleptoparasitism as a foraging strategy in anurans: Observed in the Matheran leaping-frog (*Indirana leithii*) from Maharashtra, India. *Hamadryad*. In Press.

**IMPORTANT** This current version of the accepted manuscript is a submission that has been peer reviewed and accepted for publication by *Hamadryad*. This version will have minor differences from the final publication. During the article production process, the text and other elements of this article are likely to change.

When the final article is available, the 'Accepted Manuscript' version will be removed and replaced by the final article. The date the article was first made available online will be carried over, but the year of publication will correspond to the volume/issue of inclusion.

An accepted manuscript at *Hamadryad* can be cited in text as "In Press".

**Keywords:** Amphibians, intraspecific interactions, Matheran, Ranixalidae, Western Ghats

Kleptoparasitism is a form of resource acquisition in which an individual steals an already procured resource from another individual. Although this strategy has largely been documented in the form of food theft (Clay and Rothschild 1957; Iyengar 2008), ornithologists have expanded the definition of kleptoparasitism to include the theft of other resources such as nest material (VanderWerf 1998) and nesting sites (Kappes 1997). When an individual exhibits this behaviour towards a conspecific, it is termed as intraspecific kleptoparasitism. On the contrary, when an individual exhibits this behaviour towards an individual of another species this is termed as interspecific kleptoparasitism. Individuals who find and capture their own food are crucial to the survival of the kleptoparasite. The kleptoparasites are termed as 'scroungers' who depend on the 'producers' (Nishimura 2010). Kleptoparasitism as a foraging strategy has been observed in many vertebrate taxa, including birds (Brockmann and Barnard 1979), mammals (Gorman et al. 1998; Cusack et al. 2017), fishes (Wallace and Snyder 1988), and reptiles (Platt et al. 2007). Although, kleptoparasitic behaviour for other resources like female mates and perch site has been observed in anurans (Wells 2007; Modak et al. 2018), it has been observed as a foraging strategy only in captivity (Boice and Williams 1971). I present the first observation of intraspecific kleptoparasitism as a foraging strategy in the Matheran leaping-frog, *Indirana leithii* Boulenger, 1888 from Matheran, Raigad district, Maharashtra, India at the species' type locality.

*Indirana leithii* is a species of terrestrial amphibians endemic to the Northern Western Ghats, in the states of Maharashtra and Gujarat, India (Modak et al. 2014). The type locality, Matheran, is a plateau (800 m asl), in Raigad district, Maharashtra. The vegetation on this plateau is semi-evergreen with high rainfall during the monsoon season and low temperatures relative to its surroundings. The individuals of *I. leithii* breed between June and September, and are abundant around small streams, forest floor, and human settlements in the type locality. Individuals of this species tend to congregate during the breeding season (up to 10 individuals at a site; Modak et al. 2018). Because of this aggregation during the breeding season, individuals tend to be agnostic towards each other. Males of *I. leithii* tend to exhibit kleptoparasitic behaviour towards conspecific males where they try to push out the male already in amplexus (Modak et al. 2018). Furthermore, attempted cannibalism has also been reported between individuals of the species (Kulkarni et al. 2020).

On 13<sup>th</sup> October 2019 at 23:00 h (IST), I observed intraspecific kleptoparasitism in *I. leithii* in Mathern, Raigad district, Maharashtra, India (19.0025 N, 73.2839 E). I found two adult individuals of *I. leithii* on the ground with either end of an earthworm (*Lumbricus* sp.) in their mouths (Figure 1). Both individuals attempted to pull the earthworm from each other's jaws by tugging from either end and thus causing the other individual to flip over. The flipping and tossing continued for another 5 minutes before one individual finally gave up and let the earthworm go. The victorious individual continued eating the earthworm while the other lay on its back for another 5 minutes, and then continued foraging (Figure 2). The two frogs were found adjacent to a drain with 15–20 individuals of *I. leithii*. The other individuals were actively moving and foraging.

The current observation is, to the best of my knowledge, the first record of kleptoparasitism as a foraging strategy in anurans in the wild. The choice of kleptoparasitism over other foraging

strategies depends on various factors including competitive differences among individuals, the spatial distribution of prey and host, the abundance of prey, size of the prey, handling time of the prey, and group size of host (Iyengar 2008). The observed kleptoparasitic behaviour in this species can be an outcome of the high abundance of *I. leithii* at Matheran and the aggregation of individuals at the site (Modak et al. 2018). Since the earthworm was larger in size compared to the two individuals, there was an increase in the handling time of the prey. This could be the other reason for the scrounger to exhibit this behaviour. To this end, I provide a unique account of the interactions between two individuals in the wild. My observations provide insights into anuran behaviour in an extremely rare social circumstance.

**Funding:** This short-note is an outcome of a self-funded trip to Matheran.

**Declaration:** There is no conflict of interest with other studies.

**Acknowledgement:**

I thank Mr. Shubham Soni and Dr. Seshadri K. S. for their valuable comments and inputs on the manuscript. I would also like to thank Mr. Manan Mehta and Mr. Anirudh Nippani for accompanying me in the field. I thank the editor of the journal and the anonymous reviewers for their valuable comments and suggestions.

**References:**

- Brockmann, H.J. & Barnard, C.J. (1979) Kleptoparasitism in birds. *Animal Behaviour* 27: 487–514. [https://doi.org/10.1016/0003-3472\(79\)90185-4](https://doi.org/10.1016/0003-3472(79)90185-4)
- Nishimura, K. (2010) 'Kleptoparasitism and Cannibalism', in *Encyclopedia of Animal Behavior*. Elsevier, pp. 253–258. Available at: <https://doi.org/10.1016/B978-0-08-045337-8.00279-5>
- Clay, T. & Rothschild, M. (1957) *Fleas, flukes & cuckoos; a study of bird parasites*. New York: Macmillan. <https://doi.org/10.5962/bhl.title.6413>
- Cusack, J.J., Dickman, A., Kalyahe, M., Rowcliffe, M., Carbone, C., MacDonald, D. & Coulson, T. (2017) Revealing kleptoparasitic and predatory tendencies in an African mammal community using camera traps: a comparison of spatiotemporal approaches. *Oikos* 126(6), pp. 812–822. <https://doi.org/10.1111/oik.03403>
- Gorman, M.L., Mills, M. & Raath, J. (1998) High hunting costs make African wild dogs vulnerable to kleptoparasitism by hyaenas. *Nature* 391(6666), pp. 479–481. <https://doi.org/10.1038/35131>

- Biju, S.D., Sushil, D., Anand P. & Robert, I. (2004) *Indirana leithii*, The IUCN Red List of Threatened Species 2004: e.T58311A11763222'.  
<https://doi.org/10.2305/IUCN.UK.2004.RLTS.T58311A11763222.en>
- Iyengar, E.V. (2008) Kleptoparasitic interactions throughout the animal kingdom and a re-evaluation, based on participant mobility, of the conditions promoting the evolution of kleptoparasitism: Evolution of Kleptoparasitism. *Biological Journal of the Linnean Society* 93(4), pp. 745–762. <https://doi.org/10.1111/j.1095-8312.2008.00954.x>
- Kappes, J.J. (1997) Defining cavity-associated interactions between red-cockaded woodpeckers and other cavity dependent species: interspecific competition or cavity kleptoparasitism?. *Auk* 114, pp. 778–780.
- Kulkarni, A., Nashirabadkar, A. & Patil, S. (2020) Attempted Cannibalism by a Matheran Leaping Frog, *Indirana leithii* (Ranixalidae). *Reptiles & Amphibians*, 27(2), pp. 259–260. <https://doi.org/10.17161/randa.v27i2.14262>
- Modak, N., Chunekar, H. & Padhye, A. (2018) Life History of Western Ghats endemic and threatened Anuran – Matheran leaping frog, (*Indirana leithii*) with notes on its feeding preferences. *Journal of Natural History*, 52(27–28), pp. 1745–1761.  
<https://doi.org/10.1080/00222933.2018.1488008>
- Modak, N., Padhye, A. & Dahanukar, N. (2014) Delimiting the distribution range of *Indirana leithii* (Boulenger, 1888) (Anura: Ranixalidae), an endemic threatened anuran of the Western Ghats, based on molecular and morphological analysis, *Zootaxa*, 3796(1), p. 62. <https://doi.org/10.11646/zootaxa.3796.1.3>.
- Nishimura K. (2010) Kleptoparasitism and Cannibalism, *Encyclopedia of Animal Behavior*, 253-258.
- Platt, S.G., Rainwater, T., Snider, S., Garel, A., Anderson, T. & McMurry, S. (2007) Consumption of Large Mammals by *Crocodylus moreletii*: Field Observations of Necrophagy and Interspecific Kleptoparasitism, *The Southwestern Naturalist*, 52(2), pp. 310–317.
- VanderWerf, E. (1998) Breeding Biology and Territoriality of the Hawaii Creeper, *The Condor*, 100, pp. 541–545.
- Wallace, D. & Snyder, A. (1988) Kleptoparasitism of freshwater crabs by cichlid fishes endemic to Lake Barombi Mbo, Cameroon, West Africa, *Environmental Biology of Fishes*, 22(2), pp. 155–160.
- Wells, K.D. (2007) *The Ecology and Behavior of Amphibians*. University of Chicago Press.  
<https://doi.org/10.7208/chicago/9780226893334.001.0001>



**Figure 1.** Two individuals of *Indirana leithii* with an earthworm in their mouths



**Figure 2.** One individual flipping over the other individual